

# Data Analysis of the Benefits of an Electronic Registry of Information in a Neonatal Intensive Care Unit in Greece

by Maria Skouropoulou, PhD; George Soloupis, MSc; Antonis Gounaris, MD; Antonia Charitou, MD; Petros Papasarantopoulos; Sophia L. Markantonis, PhD; Christina Golna, LL.M, MSc; and Kyriakos Souliotis, PhD

## Abstract

**Objectives:** This study assesses the results of implementation of a software program that allows for input of admission/discharge summary data (including cost) in a neonatal intensive care unit (NICU) in Greece, based on the establishment of a baseline statistical database for infants treated in a NICU and the statistical analysis of epidemiological and resource utilization data thus collected.

**Methods:** A software tool was designed, developed, and implemented between April 2004 and March 2005 in the NICU of the LITO private maternity hospital in Athens, Greece, to allow for the first time for step-by-step collection and management of summary treatment data. Data collected over this period were subsequently analyzed using defined indicators as a basis to extract results related to treatment options, treatment duration, and relative resource utilization.

**Results and discussion:** Data for 499 babies were entered in the tool and processed. Information on medical costs (e.g., mean total cost  $\pm$  SD of treatment was €310.44  $\pm$  249.17 and €6704.27  $\pm$  4079.53 for babies weighing more than 2500 g and 1000–1500 g respectively), incidence of complications or disease (e.g., 4.3 percent and 14.3 percent of study babies weighing 1,000 to 1,500 g suffered from cerebral bleeding [grade I] and bronchopulmonary dysplasia, respectively, while overall 6.0 percent had microbial infections), and medical statistics (e.g., perinatal mortality was 6.8 percent) was obtained in a quick and robust manner.

**Conclusions:** The software tool allowed for collection and analysis of data traditionally maintained in paper medical records in the NICU with greater ease and accuracy. Data codification and analysis led to significant findings at the epidemiological, medical resource utilization, and respective hospital cost levels that allowed comparisons with literature findings for the first time in Greece. The tool thus contributed to a clearer understanding of treatment practices in the NICU and set the baseline for the assessment of the impact of future interventions at the policy or hospital level.

**Keywords:** newborn, NICU, data management, statistical analysis, software

## Introduction

Birth registries maintain useful information on the number of births and perinatal morbidity and mortality in addition to providing an overview of resource utilization from the perspective of the health care system as a whole. Such data aid the definition of specific hospital procedures (both medical and administrative) and health policy initiatives that contribute to comprehensive, safe, and effective clinical treatment of newborns while at the same time allowing for robust estimation of the quality, effectiveness, and even cost-effectiveness of services provided in neonatal intensive care units (NICUs). Such implications become increasingly important in the light of the ever-growing sensitivity of

healthcare systems toward resource consumption in the hospital setting as well as the need for the provision of services of high quality to patients in need.<sup>1</sup>

In the case of premature infants treated in an intensive care unit (ICU), the availability of detailed archives is of vital importance. Two studies evaluated the causes of medication errors in NICUs and identified a broad range of errors that affect high-risk neonates.<sup>2,3</sup> Data also suggest that newborn infants are more likely to experience adverse reactions than adults and that infections are associated with high mortality among neonates.<sup>4-6</sup> The use of computerized hospital records may potentially reduce medical errors and improve the cost-effectiveness of treatments by revealing the relationship between severity of illness and resource consumption in the ICU setting.<sup>7-9</sup> Nonetheless, the implementation of a system of detailed records and the analysis of such records are both time-consuming and difficult. User-friendly software that facilitates data entry by the NICU medical and nursing staff and evaluation of economic data in parallel with statistical analysis could be a powerful tool for ensuring precise and effective interventions in this patient group. Moreover, such tools could provide useful information for parents and caregivers of hospitalized neonates.

This study recorded the outcomes of the introduction of a computerized system for registering births and treatment data in a NICU, where all relevant data were previously maintained in paper records, in order to assess the impact on the ease and accuracy of data entry as well as the availability of statistical analyses over a one-year period. The study took place in the NICU of the LITO private maternity hospital in Athens, Greece.

## Methods

In order to assess the prevalence, type, and treatment of premature neonates hospitalized in the LITO private maternity hospital, a comprehensive software program was designed that would capture all relevant data in an electronic patient record format that would in turn allow for the definition of treatment options and the calculation of total resource utilization per case.

The programming language as well as the design and development details of the software tool were approved by the scientific committee of the hospital. The software tool was developed using Visual dBase, with the minimum hardware requirements as follows: CPU: Celeron; RAM: 256 MB; HD: 500 MB free; operating system: Windows 98, 2000, or XP.

Demographic and clinical data of patients admitted into the NICU of the LITO maternity hospital between April 1, 2004, and March 31, 2005, were loaded into the software. Data were inputted and analyzed for 499 newborn children, and the cost for their hospitalization was calculated. First-time users of the program were timed while inputting data for 100 newborns at the beginning and end of the one-year study period (at which point they were characterized as experienced users), and the time required to input data was compared. Data entry was closely monitored and assessed by computer scientists so that further improvements to the program could be made in accordance with feedback from the clinicians.

For each patient record, the software collected

- (a) demographical and epidemiological data related to the mother;
- (b) epidemiological data related to the premature neonate (prevalence of condition or conditions, treatment requirements, etc.); and
- (c) resource utilization data (treatment and hospitalization cost), which were validated against data maintained in separate records at the finance department.

The tool consisted of basic, easy-to-follow steps that allowed for the capturing of relevant epidemiological and resource utilization data:

- (a) The application requested a security access code. Each user had unique access codes (username and password) to ensure privacy of data.
- (b) The user entered data related to the mother, including information about her clinical and personal history (see Figure 1). The user selected options, where appropriate, from drop-down menus or created new entries.

- (c) The user entered data on the diagnoses and health conditions of the infant. The specific diagnosis or diagnoses could be selected from the archive of diagnoses available from the program's database (see Figure 2). Each selected diagnosis was automatically saved.
- (d) Information pertaining to a neonatal group of potential interest could be selected by pressing the "summarized printing" button. When the "analytical printing" option was selected, a detailed report including individual data from each neonate could also be printed. All relevant details could be summarized in a comprehensive form (see Figure 3).
- (e) Resource utilization data were entered in the relevant fields. These fields were defined in cooperation with the finance department and reflected the total cost of treatment that was charged to the family or payer (insurance fund) for the duration of the treatment of the infant. The cost was broken down to specific cost centers in order to fully reflect resource utilization according to the clinical condition of each infant.

Statistical analyses used unpaired *t*-tests for the comparison of means and  $\chi^2$  tests for the comparison of percentages. Spearman correlation coefficients were calculated for selected variables. Significance was defined as  $p < .05$ .

## Results

Over the one-year study period, the data assembled and analyzed through the use of the software tool indicated a perinatal mortality rate of 6.8 percent. The infants in the study had been treated at normal (Level I), intermediate (Level II), and intensive-care nursery (Level III) levels as shown in Table 1. The demographic characteristics of the mothers were also entered into the software (see Table 2).

Twenty-nine percent of the babies were born at a gestational age of greater than 37 weeks. Table 3 depicts the number of infants in each of the four different categories according to their birth weights. For each weight category, the mean length of stay, the mean cost, and the most important complications were recorded (see Table 4).

Fourteen percent of the mothers had undergone in vitro fertilization. This corresponds to 72 newborn infants, of whom 16.6 percent (12 newborns) were singletons, 61.1 percent (44 newborns) were twins, and 22.3 percent (16 newborns) were triplets or quadruplets. Their mean birth weight was  $1924 \pm 481.8$  g, and their mean gestational age was  $34 \pm 3$  weeks. Level III nursery care was provided for 62.5 percent of the babies, and the rest received lower levels of nursery care.

As expected, both the gestational age at delivery and the birth weight decreased as the number of fetuses increased, that is, with twin or greater gestation. The cost of treatment was high when the baby suffered respiratory complications. In the category of birth weights from 1,000 to 1,499 g at Level III, bronchopulmonary dysplasia inflicted 14.3 percent of the babies.

In the category of birth weight less than 1,000 g, there were only 13 infants, eight of whom died between the first and fourth day, while the other five were transferred to other state hospitals because of financial considerations. The duration of stay and the mean cost for babies in this category apply only to the babies that passed away while in the LITO intensive care unit.

Bronchopulmonary dysplasia and cerebral bleeding inflicted 14.3 percent and 4.3 percent of babies weighing between 1,000 and 1,499 g, while the respective percentages reported by Lemons et al. were 15 percent and 14.5 percent.<sup>10</sup>

The data also confirmed that twin or greater gestation, diabetes, and rupture of membranes for more than 18 hours are significantly correlated with the length of stay (see Table 5).

No significant correlation was found between birth weight and maternal smoking,  $p = .778$ . This, however, may not be an accurate finding as only 40 mothers (8 percent) smoked during pregnancy, making the sample too small to allow for a valid conclusion. The mean weight of the babies whose mothers smoked before and during pregnancy was  $2629 \pm 651.3$  g, their mean gestational age was  $37 \pm 3$  weeks (range 31–40 weeks), and the mean duration of their stay in the ICU was  $8 \pm 5.3$  days.

Finally, data were collected on costs incurred for each category of babies and compared with the available literature.<sup>11,12</sup> The mean total cost of treatment was calculated at  $\text{€}310.44 \pm 249.17$  and  $\text{€}6704.27 \pm 4079.53$  for babies weighing more than 2,500 g and 1,000–1,500 g, respectively. The study

also concluded that when a baby's weight is under 1,500 g, the cost for treatment and care is 60 percent higher than when the baby weighs between 1,500 and 2,499 g and 130 percent higher than the cost for a baby weighing more than 2,500 g.

## Discussion and Conclusion

The importance of computerized data management to increase safety and efficacy in an ICU for premature neonates has been fully realized over the past decades. The electronic documentation of several procedures for neonates, such as parenteral nutrition in the ICU, has been referred to in the literature.<sup>13,14</sup> The establishment of a monitoring system allows for outcome research as well as for management of information.<sup>15</sup> Calculated yearly statistical outcomes have been shown to be very important for the protection of future neonates and for the optimization of the function of an ICU. The availability of this information allows clinicians to minimize mistakes and reevaluate current clinical practices. Such analysis has become increasingly important as numbers of live births registered in most industrialized nations have remained relatively stable in recent years, but the incidence of preterm birth and low birth weight is increasing.<sup>16</sup>

Over the past two years a software program that combines fast report generation and a capacity for simple statistical analyses was developed and used to collect, store, and analyze data of newborns treated in all three levels of the intensive care unit of the LITO maternity hospital. Over this period the software tool enabled the breakdown of data by types of patients, different diagnoses, operations performed, and any medical or surgical complications incurred.

Data collected and managed through the tool revealed that babies weighing between 1,000 and 1,500 g who were treated in the NICU of the LITO maternity hospital suffered fewer complications than would be expected based on reports in the literature. Findings on twin or greater gestation, diabetes, and rupture of membranes of greater than 18 hours are in line with previous studies.<sup>17</sup> The length of stay in the NICU was shorter than that reported elsewhere, where the mean length of stay was 40 days.<sup>18</sup> The mean cost of hospitalization was lower than the one reported in the United States, where the mean cost amounted to USD 1,100, and approximately equal to costs estimated from a meta-analysis of 15 studies in the United Kingdom (mean cost of GBP 225).<sup>19,20</sup>

Users credited the tool with harmonizing and codifying the documentation of various diseases and treatment pathways of premature infants. The software tool thus simplified the registration process for all employees involved, as they had to select from previously defined options. It also allowed for automatic archiving of data as well as for automatic retrieval of data to prepare statistical reports. Complete and accurate summaries of all admissions could be prepared at the time of a baby's discharge or upon request. At the service provision level, the program offered an overview of the efficacy of clinical practices, as treatment pathways could be recorded in full and compared against best-practice guidelines. The production of periodic reports of each department's and each practitioner's activities offered a clearer understanding of currently established practices and initiated discussions at the administrative level to harmonize practices, define benchmarks for service quality and cost, and optimize efficiency.

This study evaluated the introduction for the first time in Greece of an electronic registry of information for premature infants. The software tool thus developed and implemented has the potential to assist all caregivers in NICU settings, as it allows for harmonized, codified collection of data on the treatment pathway of each infant as well as the systematic and consistent production of activity reports. Data thus stored and analyzed can be a useful tool for parents and can help hospital administrators, insurance companies, and government agencies to design health policy interventions that comply with best-practice guidelines.

Maria Skouroliaou, PhD, clinical pharmacist, is a lecturer of parenteral and enteral nutrition at Harokopion University, Athens, Greece

George Soloupis, MSc, pharmacist, is a postgraduate student at the School of Pharmacy of the University of Athens, Greece

Antonis Gounaris, MD, PhD, is an assistant professor of neonatology and director of the NICU at the General Hospital of Piraeus, Agios Panteleimon, Greece

Antonia Charitou, MD, is a neonatologist and director of the NICU at the Lito Maternity Hospital in Athens, Greece

Petros Papasarantopoulos, is a pharmacist at Business Information Support in Athens, Greece

Sophia L. Markantonis, PhD, pharmacist, is an associate professor at the School of Pharmacy of the University of Athens, Greece

Christina Golna, LL.M, MSc, lawyer, is a Health Policy and Patient Access Manager at Roche (Hellas) SA

Kyriakos Souliotis, PhD, is a lecturer of health policy at the Department of Social and Education Policy at the University of Peloponnese, Greece, and a chief planning officer at the Health Fund of the National Bank of Greece

## Notes

1. Golna, C., and Souliotis, K. "Ageing and Health Policy." In K. Souliotis (Editor), *Health Policy and Economics*. Athens, Greece: Papazisis, 2006, pp. 21–44 (in Greek).
2. Suresh, G., Horbar, J. D., Plsek, P., et al. "Voluntary Anonymous Reporting of Medical Errors for Neonatal Intensive Care." *Pediatrics* 113, no. 6 (2004): 1609–1618.
3. Vincer, M. J., J. M. Murray, A. Yuill, et al. "Drug Errors and Incidents in a Neonatal Intensive Care Unit." *American Journal of Diseases of Children* 143 (1989): 737–740.
4. Van den Anker, J. N. "Managing Drugs Safely." *Seminars in Fetal and Neonatal Medicine* 10 (2005): 73–81.
5. Storgion, S. A., R. J. Beck, and R. J. Leggiadro. "Frequency and Outcome of Infectious-disease Admissions to a Paediatric Intensive-care Unit." *Southern Medical Journal* 87, no. 11 (1994): 1121–1124.
6. Arafá, M. A., and M. A. Alshehri. "Predictors of Neonatal Mortality in the Intensive Care Unit in Abha, Saudi Arabia." *Saudi Medical Journal* 24, no. 12 (2003): 1374–1376.
7. Sado, A. S. "Electronic Medical Record in the Intensive Care Unit." *Critical Care Clinics* 15 (1999): 499–522.
8. Lowe, W. W., T. A. Ciszek, and K. J. Gallaher. 1992. "Comprehensive Computerized Neonatal Intensive Care Unit Data System Including Real-time Computer Generated Daily Progress Notes." *Pediatrics* 89, no. 1 (1992): 62–66.
9. Martinot, A., S. Leteurtre, B. Grandbastien, et al. "Patient Characteristics and Resource Utilization in Nine French Paediatric Intensive Care Units." *Archives de pédiatrie* 4, no. 8 (1997): 730–736.
10. Lemons, J. A., C. R. Bauer, W. Oh, et al. "Very Low Birth Weight Outcomes of the National Institute of Child Health and Human Development Neonatal Research Network, January 1995 Through December 1996." *Pediatrics* 107, no. 1 (2001): E1.
11. Phibbs, C. S., R. L. Williams, and R. H. Phibbs. "Newborn Risk Factors and Costs of Neonatal Intensive Care." *Pediatrics* 68 (1981): 313–321.
12. Doyle, L. W., L. J. Murton, and W. H. Kitchen. "Increasing the Survival of Extremely Immature (24 to 28-weeks gestation) Infants—at what cost?" *Medical Journal of Australia* 150 (1989): 558–568.
13. Schloerb, P. R. "Electronic Parenteral and Enteral Nutrition." *Journal of Parenteral and Enteral Nutrition* 24 (2000): 23–29.
14. Philipp, F., M. Jantke, E. Finkeissen, et al. "Introducing DoT-U2—an XML-based Knowledge Supported Checklist Software for Documentation of a Newborn Clinical Screening Examination." *Computer Methods and Programs in Biomedicine* 77, no. 2 (2005): 115–120.
15. Tellen, S. "Developing a State Outcome Monitoring System in Public Health Maternal and Child Health." *Journal of Medical Systems* 23, no. 3 (1999): 227–238.
16. Macfarlane, A., and M. Mugford. "The Outcome of Pregnancy—What Can We Learn from Official Statistics?" *Nursing Times* 80, no. 20 (1984, May 16–22): 42–44.
17. Ross, M. G., C. A. Downey, R. Bemis-Heys, et al. "Prediction by Maternal Risk Factors of Neonatal Intensive Care Admissions: Evaluation of >59,000 Women in National Managed Care Programs." *American Journal of Obstetrics and Gynecology* 181 (1999): 835–842.
18. Rogowski, J. "Using Economic Information in a Quality Improvement Collaborative." *Pediatrics* 111 (2003): 411–418.
19. Rogowski, J. "Using Economic Information in a Quality Improvement Collaborative."
20. Petrou, S. "Economic Consequences of Preterm Birth and Low Birthweight." *British Journal of Obstetrics and Gynaecology* 110, no. 20 (2003): 17–23.

**Table 1**

Level of Nursery Care for Infants

Level of nursery	<i>N</i>	Percentage
Normal (Level I)	268	53.7
Intermediate (Level II)	90	18.0
Intensive care (Level III)	141	28.3

**Table 2**Characteristics of Study Population ( $n = 499$ )

Characteristic	<i>N</i>	Percentage
<b>Gender</b>		
Male	230	46.1
Female	269	53.9
<b><i>Delivery</i></b>		
Vaginal	162	32.5
Cesarean	337	67.5
<b><i>Gestation</i></b>		
Singleton	378	75.8
Twin	105	21.0
Triplet	15	3.0
Quadruplet or above	1	0.2



**Table 3**

Birth Weight Categories of Babies

Birth weight (g)	<i>N</i>	Percentage
>2500	245	49.1
1500–2499	212	42.5
1000–1499	29	5.8
<1000	13	2.6

**Table 4**

Length of Stay, Cost, and Clinical Complications

Birth weight (g)	Level of nursery	<i>N</i>	Duration of stay (days) (mean ± SD)	Respiratory diseases (%)	Microbial infections (%)	Cerebral bleeding (%)	Total cost (€) (mean ± SD)
>2500	Level I	138	4 ± 3.5	5.07	0.7	0	310.4 ± 249.2
	Level II	34	6.2 ± 2.7	20.6	8.8	5.9	1176.8 ± 670.2
	Level III	73	9.0 ± 4.3	80.8	5.5	0	2945.4 ± 1990.1
1500–2499	Level I	104	5.5 ± 2.9	1.9	0.0	0.0	947.6 ± 760.1
	Level II	32	8.6 ± 4.3	21.4	4.8	0.0	1596.0 ± 555.6
	Level III	76	13.5 ± 7.9	78.9	7.9	0.0	4201.3 ± 3334.6
1000–1499	Level I	0	-	-	-	-	-
	Level II	4	8.5 ± 2.1	0	0	0	-
	Level III	25	21.7 ± 17.1	78.3	8.0	4.3	6704.3 ± 4079.5
<1000	Level III	13	2.2 ± 1.5	76.9	30.8	23.1	1647.6 ± 603.8

*Note:* A dash (-) indicates that no babies were within a category.

**Table 5**

Maternal Factors Associated with Length of Stay

	<i>Length of stay in days (mean ± SD)</i>	<i>p</i>
Diabetes		
Yes	10.4 ± 9.4	0.049
No	7.8 ± 6.9	
Rupture of membranes >18h		
Yes	11.6 ± 8.3	0.034
No	7.8 ± 6.9	
Twin or greater gestation		
Yes	10.3 ± 9.5	0.002
No	7.4 ± 6.2	
Abruptio placentae		
Yes	12.9 ± 13.2	0.154
No	7.9 ± 7.9	

**Figure 1**

Form Providing Summary Epidemiological Information on the Mother

The screenshot shows a Visual dBASE application window titled 'Visual dBASE' with a menu bar containing 'FILE', 'EDIT', 'STATISTIC', and 'UTILITIES'. Inside the window, there is a sub-window titled 'B.I.S' containing a form titled 'NEONATES FILE page 1/2'. The form is organized into several sections:

- Header Section:** Includes fields for 'CODE', 'NEONATE'S SURNAME' (with a search icon), 'SEX', 'BIRTH DATE' (formatted as / /), 'BIRTH WEIGHT', and 'GESTAT'.
- Medical History Section:** Contains dropdown menus for 'PEDIATRICIAN', 'TYPE OF DELIVERY', and 'MULTIPLE GESTATION'.
- Maternal Information Section:** Includes fields for 'NEONATE UNIT', 'MOTHER'S SURNAME', and 'MOTHER'S NAME'.
- Contact and Status Section:** Includes fields for 'ADDRESS', 'PHONES', 'GYNECOLOGIST', 'FAMILY STATUS', 'BORN', and 'MATERNAL PROBLEMS (1)'. There are also dropdown menus for 'MATERNAL PROBLEMS (2)', 'MATERNAL PROBLEMS (3)', and 'PRENATAL STEROID ADMINISTRATION'.
- Remarks Section:** A text area labeled 'REMARKS' with a 'memo' button.
- Navigation and Action Buttons:** A row of buttons at the bottom includes 'First', 'Previous', 'Next', 'Last', 'New' (with a green checkmark), 'Delete' (with a red X), 'Exit' (with a computer icon), and 'Next page' (with a blue arrow).

The status bar at the bottom of the window displays 'lsex.dbf', 'Rec 1/1', and 'Ins'.

**Figure 2**

Form Summarizing Information on the Health State of the Infant

The screenshot shows a Visual dBASE application window titled "Visual dBASE" with a menu bar (FILE, EDIT, STATISTIC, UTILITIES) and a toolbar. The main window displays a form titled "NEONATES FILE" on "page 2/2".

The form is organized into three columns:

- Column 1 (Left):** Contains fields for "NEONATAL RESUSCITATION", "RESPIRATORY DISEASES", "RESPIRAT. TREATMENT" (with a "DAYS" spinner), "CONGENITAL INFECTIONS", "CEREBRAL BLEEDING", "CONGENITAL METABOLIC DISORDERS", "NUTRITION", "NEUROLOGICAL CONDITION AT EXIT", and "READMISSION".
- Column 2 (Middle):** Contains fields for "APGAR" (with spinners for 1', 5', and 10'), "RESPIRATORY DISEASES TREATMENT", "ADMINISTRATION SURFACTANT", "MICROBIAL INFECTIONS", "CIRCULATORY DISEASES", "VISION DISORDERS", "OTHER DIAGNOSES", "EXIT", and "FOLLOW UP".
- Column 3 (Right):** Contains fields for "NEONATAL JAUNDICE", "NASAL CPAP", "MECHANICAL VENTILATION COMPLICATIONS", "PERINATAL ASPHYXIA (ANOXIA)", "CONGENITAL DISORDERS", "HEARING DISORDERS", "SURGICAL", "DISCHARGE AGE" (with a "DAYS" spinner), and "OTHER".

At the top of the form, there are search fields for "NEONATE'S SURNAME", "SEX", "MOTHER'S SURNAME", and "MOTHER'S NAME".

At the bottom of the form, there is a control bar with buttons: "First", "Previous", "Next", "Last", "New" (with a green checkmark), "Delete" (with a red X), and "Exit" (with a computer icon). The status bar at the very bottom shows "lserx.dbf", "Rec 1/1", and "Ins".

**Figure 3**

Form Summarizing Information Provided for Each Mother and Neonate

<b>Address</b>				<b>Phone No 1</b>			
<b>Gynecologist</b>				<b>Phone No 2</b>			
<b>Type of Room</b>		Single Bed Room		Two-Bed Room		Lux Room	
<b>Born</b>	First Born		Second Born		Third Born		Fourth Born & Over
<b>Sex of the Baby</b>				<b>Date of Birth</b>			
<b>Birth Weight</b>				<b>Gestational Age</b>		<b>Pediatrician</b>	
<b>Type of Delivery</b>				<b>Multiple Gestation</b>		<b>Neonate Unit</b>	
EMERGENCY CESAREAN SECTION (CS) SECONDARY TO PROLAPSED CORD				Single Gestation		Admission at 1 <sup>ST</sup> level	
EMERGENCY CS SECONDARY TO FETAL DISTRESS				Twin Gestation		Admission at 2 <sup>ND</sup> level	
EMERGENCY CS SECONDARY TO LABOR DYSTOCIA				Triple Gestation		Admission at 3 <sup>ND</sup> level	
EMERGENCY CS SECONDARY TO PLACENTA ABRUPTION				Multiple Gestation		Observations	
EMERGENCY CS SECONDARY TO MATERNAL INDICATION							
Elective CS							
Normal – Breech Presentation							
<b>Maternal Problems</b>							
Bleeding During Gestation		Amniocentesis		Placenta Abruption			
Diabetes		In Vitro Fertilization (IVF)		Smoker			
Maternal Infection		Drugs – Narcotics		Oligamnio			
Pathological Biophysical Profile		Prenatal Ultrasound (US)		Rupture of Membranes <18H			
Premature Rupture of Membranes >18H		Congenital Infection		Toxemia			
Hydramnio		Observation: write down up to three problems					
<b>Prenatal Steroid Administration</b>		One Dose		Two Doses			
		Three Doses					
<b>Neonatal Resuscitation</b>		<b>Neonatal Jaundice</b>		<b>Respiratory Diseases</b>			
No Need		Jaundice-No Need for Treatment		Respiratory Distress Syndrome			
Respiratory Mask (Ambu)		Unknown Cause – Phototherapy		Transient Tachypnea of newborn			

Respiratory Mask (Ambu)- Intubation		Unknown Cause - Exchange Transfusion		Spontaneous Pneumothorax	
Respiratory Mask (Ambu) Intubation Cardiac Massage		Rhesus Incompatibility- Phototherapy		Meconium Aspiration	
Respiratory Mask (Ambu)- Intubation-Cardiac Massage Adrenaline (Epinephrine)		Rhesus Incompatibility - Exchange Transfusion		Congenital Pneumonia	
OTHER		ABO Incompatibility- Phototherapy		Persistent Fetal Circulation	
		ABO Incompatibility - Exchange Transfusion			
<b>Perinatal Asphyxia (Anoxia)</b>					
Mild APGAR Score <7 at 1'					
Medium APGAR score 4-7 at 5'					
Severe APGAR Score <4 at 5'					
Seizures – Convulsions					
<b>Respiratory Disease Treatment</b>					
Diffuse F <sub>1</sub> O <sub>2</sub>		nCPAP+PTV		Diffuse F <sub>1</sub> O <sub>2</sub> +nCPAP+PTV	
Diffuse F <sub>1</sub> O <sub>2</sub> +HOOD		nCPAP+PTV+HFO		Diffuse F <sub>1</sub> O <sub>2</sub> +nCPAP+PTV+HFO	
n.CPAP		nCPAP+PTV+HFO+NO		Diffuse F <sub>1</sub> O <sub>2</sub> +nCPAP+PTV+HFO+NO	
SIMV- PTV		PTV+HFO		Other	
<b>Duration of Respiratory Treatment (Days)</b>					
<b>Nasal CPAP</b>					
		<b>Cerebral Bleeding</b>		<b>Other Diagnoses</b>	
Ventilator Apparatus		First-Grade Intraventricular Hemorrhage (IVH)		Simple Observation	
Bubble CPAP		Second-Grade IVH		Simple Prematurity	
<b>Administration Surfactant</b>		Third-Grade IVH		Newborn of Diabetic Mother	
Administration CUROSURF 1 Time		Fourth-Grade IVH		Congenital Laryngeal Inspiration	
Administration CUROSURF 2 Times		Periventricular Leucomalacia (PVL)		Cardiorespiratory Failure	
Administration CUROSURF 3 Times		Methemorrhagic Hydrocephalus		Heart Failure	
Administration CUROSURF >3 Times		Intracerebellar Hemorrhage		Peritonitis	
Administration SURVANTA 1 Time		Subarachnoid Hemorrhage		DIC	
Administration SURVANTA 2 Times		Subdural Hemorrhage		IUGR	

Administration SURVANTA 3 Times		GOR-Gastroesophageal Reflux	
Administration SURVANTA >3	<b>Circulatory Diseases</b>	<b>Nutrition</b>	
	Acyanotic Congenital- Open Ductus Arteriosus- Indocid	Oral with Maternal Milk	
<b>Mechanical Ventilation Complications</b>	Acyanotic Congenital- Open Ductus Arteriosus- No Indocid	Oral with Maternal Milk + Fortifier	
Bronchopulmonary Dysplasia-Mild	Acyanotic Congenital- Open Ductus Arteriosus- Ligation	Oral with Special Formula	
Bronchopulmonary Dysplasia-Moderate	Acyanotic Congenital-Aortic Stenosis	Oral with Maternal and Normal Milk	
Bronchopulmonary Dysplasia-Severe	Acyanotic Congenital- Atrioventricular Communication- AV. Canal	Oral with Maternal and Special Formula	
Pneumothorax	Acyanotic Congenital – Ventricular Septal Defect (VSD)	Parenteral for <7 Days	
Interstitial Emphysema	Acyanotic Congenital – Atrial Septal Defect (ASD)	Parenteral for 7 – 28 Days	
Pneumomediastinum	Acyanotic Congenital–Peripheral Pulmonary Stenosis	Parenteral for >28 Days	
Pulmonary Hemorrhage	Acyanotic Congenital-Fallot Tetralogy		
Subcutaneous Emphysema	Acyanotic Congenital Ligation PDA	<b>Surgical</b>	
<b>Congenital Infections</b>	Cardiac Arrhythmia	Congenital Hydronephrosis	
CMV	Cyanotic Congenital-Tricuspid Ebstein Disorder	Cryptorchidism	
Coxsackie B (Enteroviruses)	Cyanotic Congenital-Arterial Caudex	Harelip-Cleft Palate	
HIV	Cyanotic Congenital-Pulmonary Atresia	Pyloric Stenosis	
PARVOVIRUS 8-19	Cyanotic Congenital-Tricuspid Atresia	Oesophageal Atresia	
Chickenpox	Cyanotic Congenital Transposition of Great Vessels	Ovary Bladder	
Viral Infection Check-Up	Cyanotic Congenital Single Ventricle	Atresia Gut	
Herpes Simplex	Myocardiopathy	Diaphragmatic Atresia	
Rubella			
Hepatitis		<b>Neurological Condition at Exit</b>	
Mycoplasma		Fine	
Lyme Disease	<b>Congenital Disorders</b>	Suspicious	
Syphilis	Unknown Syndrome	Pathologic	
Toxoplasmosis	Known Syndrome		
Chlamydia	Multiple Congenital Disorders	<b>Exit</b>	
	Congenital Encephalocele	Discharge Home	
	Congenital Meningocele	Discharge To Another Hospital	
<b>Microbial Infections</b>			
Microbial Infection Check-Up	Congenital Myelomeningocele	Transfer To Morgue (Death)-Autopsy Consent	
Necrotizing Enterocolitis (NEC)	Congenital Hydrocephalus	Transfer To Morgue (Death)-No Autopsy Consent Accept	



Suspected Necrotizing Enterocolitis (NEC)	Trisomy 13-15		
Urine Infection	Trisomy 18		
Early Onset Meningitis (- ENY)	Trisomy 21		
Early Onset** Meningitis (+ ENY)	Chromosomal Disorders		
Early Onset E coli Septicemia			
Early Onset Klebsiella Septicemia	<b>CONGENITAL METABOLIC DISORDERS</b>	<b>Readmission</b>	
Early Onset Serratia Septicemia	Congenital Hypothyroidism	Readmission Cause	
Early Onset Staph COAG (-) Septicemia	Congenital Adrenal Hyperplasia	Age (DAYS)	
Early Onset Staph COAG (+) Septicemia	Galactosemia	Transfer To Morgue (Death)-No Autopsy Consent Accept	
Early Onset Strep Type B Septicemia	Ambiguous Genitalia	Transfer To Morgue (Death)-Autopsy Consent	
Early Onset Septicemia From Other Microbes	Other Metabolic Disturbance	Discharge Home	
Early Onset Fungal Septicemia		Transfer To Ward	
Septic Arthritis	<b>Vision Disorders</b>	Hospital Stay (During Readmission)	
Late Onset* Meningitis with Negative Cerebral + Spinal Fluid (-CSF)	Normal		
Late Onset Meningitis with Positive Cerebral + Spinal Fluid (+ CSF)	ROP Stage I	<b>Discharge age( days)</b>	
Late Onset E Coli Septicemia	ROP Stage II	<b>Total Cost (€)</b>	0.00
Late Onset Klebsiella Septicemia	ROP Stage III Without Photocoagulation		
Late Onset Serratia Septicemia	ROP Stage III With Photocoagulation	<b>Follow Up</b>	
Late Onset Staph COAG (-) Septicemia	ROP Stage IV Without Photocoagulation	Severe Mental Deficit Hysteresis	
Late Onset Staph COAG (+) Septicemia	ROP stage IV With Photocoagulation	Severe Hemiplegia	
Late Onset Strep Type B Septicemia	Congenital Glaucoma	Severe Monoplegia	
Late Onset Septicemia from Other Microbes	Congenital Eye Disorders	Severe Tetraplegia	
Late Onset Fungal Septicemia	Congenital Cataract	Severe Triplegia	
		Severe Bilateral Blindness	
	<b>Hearing Disorders</b>	Moderate Deafness	
	Normal	Unilateral Blindness	
	Bilateral Absence of Acoustic Dynamics	Moderate Dysfunction – Lenient Monoplegia Diplegia	
	Unilateral Absence of Acoustic Dynamics	Moderate Dysfunction - Mild Neuro-Developmental Disturbances	

	Complete Deafness	Normal Child	
--	-------------------	--------------	--

\* late onset >7 days

\*\*early onset <3 days